

## **DIRECTIONS FOR USE**

### **General:**

For indoor use only. DO NOT apply directly to livestock and poultry. Avoid the treatment of extremely dusty areas. Do not apply product to treated supports, pieces of timber, sheets of cardboard or fabric which will be placed in areas subject to washing/sluing. Place painted supports or bait stations in areas inaccessible to food producing animals, children and pets, and where foodstuffs will not become contaminated. Best results will be achieved when small amounts of the bait are applied in many locations, concentrating around the preferred resting sites of flies.

### **APPLICATION:**

#### **DIRECT Application:**

200 g bait 100 m<sup>2</sup> of floor area. Directly place granules in bait stations at an application rate of 200 g bait per 100 m<sup>2</sup> of floor area. The granules can be directly placed within bait stations, away from pathways and/or manure and out of reach from non-target animals.

#### **PAINT-ON Application:**

200 g bait (diluted in 150 ml water) onto 1 m<sup>2</sup> discrete surface, such as cardboard sheets, timber sheets or fabric, to cover a 100 m<sup>2</sup> of floor area.

#### **Preparation:**

For the mixing/loading and the product application step, the applicator must wear disposable clothes (e.g. paper smocks, aprons, overalls) to avoid emissions to the sewer system due to washing contaminated clothing. The area where mixing/loading/application to supports, discrete surfaces, cardboard sheets, hang-boards or cards (animal husbandry\*) take place, must be covered with a disposable plastic sheet in, order to avoid contamination of adjacent surfaces and floor. Leave an untreated strip in order to handle the treated timber/sheets of card: **DO NOT touch/contact the treated area of the hang-board.**

\* In animal husbandry facilities with bare soil, application can only be made by painting onto cards.

### **Application**

200 g of bait is to be mixed with 150 mL of warm water to form a liquid paste. This is applied to 1 m<sup>2</sup> discrete surface, such as cardboard sheets, to cover a 100 m<sup>2</sup> floor area. Using a paint brush or roller, the solution is applied as stripes or spots onto pieces of timber, or sheets of cardboard or fabric which are hung or attached in areas where flies tend to aggregate. These areas may be walls, on surfaces of pillars and around windows or other structures. Treated boards should be placed in as many places as possible where flies tend to congregate (without exceeding 200 g of QuickBait per 100 m<sup>2</sup> floor/ground space).

### **Cleaning up:**

DO NOT discharge any waste product or washing water from treatment equipment to drains. Equipment used in connection with the paint-on application (e.g. brushes, roller) must not be cleaned after use. All contaminated equipment may be re-used without cleaning (if possible) and always be disposed of according to local regulations, (with no discharge into waste water).

### **Duration to effect:**

QuickBait can provide long lasting housefly control in rural hygiene and waste management facilities. However, this is likely to be best achieved if the treatment is complemented by good hygiene practice. Noticeable reduction of house fly numbers can be observed from day 1 and house fly control can be expected  $\geq 3$  weeks and can last up to 8 weeks under standard conditions. QuickBait lasts for up to 8 weeks. The minimum retreatment period is 21 days. QuickBait should not be used more than 8 times in any 12 month period.

GB05962454H - A5 - ARTICLE : 05814250

## **FIRST AID**

**General:** Move out of dangerous area. Place and transport victim in stable position (lying sideways). Remove contaminated clothing immediately and dispose of safely.

**Skin Contact:** Wash off thoroughly with plenty of soap and water, if available with polyethyleneglycol 400, subsequently rinse with water.

**Eye Contact:** Rinse immediately with plenty of water, also under the eyelids, for at least 15 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Get medical attention if irritation develops and persists.

**Ingestion:** Rinse mouth. Do not give fluids or induce vomiting. Call physician or poison control centre immediately.

**Treatment:** Treat symptomatically.

**Monitor:** Respiratory and cardiac functions.

**There is no specific antidote.**

**Poison Information for Ireland: For information or to report a poisoning incident contact The National Poisons Information Centre, Beaumont Hospital, Dublin (01-8092166), retain the label for reference.**

## **RESISTANCE MANAGEMENT:**

This product should not be used continuously against houseflies in intensive or controlled environment animal houses as this could cause failure due to insecticide resistance.

Imidacloprid is the active component of QuickBayt, which belongs to the class of neonicotinoid insecticides having a different mode of action to other classes of insecticide, such as pyrethroids or organophosphates.

The use pattern as paint-on bait ensures that most of the animal house is not treated thereby reducing the likelihood of contacting a sub lethal deposit. Nevertheless, to minimize the chances of resistance developing in the future, it is advisable to avoid using QuickBayt, exclusively and continuously as the sole agent for fly control.

QuickBayt should be used as one component of an integrated pest management program which features products from alternative chemical classes and non-chemical means (e.g. fly screens, sanitary practices).



# QuickBayt<sup>®</sup>

## INSECTICIDE

- ✓ For use only as an insecticide (PT18).
- ✓ For use only by professional operators.
- ✓ For indoor use only.

### Safety information

**QUICKBAYT** - Contains 0.5% w/w Imidacloprid & 0.1% Muscalure.



**WARNING**

**Very toxic to aquatic life with long lasting effects.**

Collect spillage.

Avoid release to the environment.

Contains Cis-tricos-9-ene. May produce and allergic reaction.

Dispose of contents / container to a licensed hazardous-waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste.

**To avoid risks to human health and the environment, comply with the instructions for use.**

UK-2017-1090, IE/BPA 70269

# 2 kg e



Paint on fly control



QuickBayt is an insecticidal water dispersible granular (WG/GR) bait for the control of flies (*Musca domestica*) for use in animal units or agricultural buildings (e.g. broiler houses, livestock barns, caged layer hen houses and small animal husbandry e.g. kennels etc.) as well as sheltered waste management facilities (only facilities which are protected from the weather, like waste sorting units). QuickBayt contains 0.5% Imidacloprid and 0.1% Muscalure for direct application or for mixing with water and spreading as a paint on treatment.

For use indoors:

- Animal houses/shelters
- Sheltered waste management sites

Fast knockdown and long lasting control (population reduction observable on day one of treatment and lasts up to 8 weeks). QuickBayt is an insecticidal water dispersible granular (WG/GR) bait containing 0.5% Imidacloprid and 0.1% Muscalure. The appeal of the bait to flies derives from its sugar base, the unique mix of flavoring and a pheromone (Muscalure®) together with its red colour. QuickBayt also contains a bittering agent, BITREX™, to deter ingestion by people, pets and food producing animals.

#### DIRECTIONS FOR USE

##### General:

For indoor use only. DO NOT apply directly to livestock and poultry. Avoid the treatment of extremely dusty areas. Do not apply product to treated supports, pieces of timber, sheets of cardboard or fabric which will be placed in areas subject to washing/slaicing. Place painted supports or bait stations in areas inaccessible to food producing animals, children and pets, and where foodstuffs will not become contaminated. Best results will be achieved when small amounts of the bait are applied in many locations, concentrating around the preferred resting sites of flies.

##### APPLICATION:

###### DIRECT Application:

200 g bait 100 m<sup>2</sup> of floor area. Directly place granules in bait stations at an application rate of 200 g bait per 100 m<sup>2</sup> of floor area. The granules can be directly placed within bait stations, away from pathways and/or manure and out of reach from non-target animals.

###### PAINT-ON Application:

200 g bait (diluted in 150 ml water) onto 1 m<sup>2</sup> discrete surface, such as cardboard sheets, timber sheets or fabric, to cover a 100 m<sup>2</sup> of floor area.

##### Preparation:

For the mixing/loading and the product application step, the applicator must wear disposable clothes (e.g. paper smocks, aprons, overalls) to avoid emissions to the sewer system due to washing contaminated clothing. The area where mixing/loading/application to supports, discrete surfaces, cardboard sheets, hang-boards or cards (animal husbandry\*) take place, must be covered with a disposable plastic sheet in, order to avoid contamination of adjacent surfaces and floor. Leave an untreated strip in order to handle the treated timber/sheets of card: DO NOT touch/contact the treated area of the hang-board.

\* In animal husbandry facilities with bare soil, application can only be made by painting onto cards.

##### Application

200 g of bait is to be mixed with 150 mL of warm water to form a liquid paste. This is applied to 1m<sup>2</sup> discrete surface, such as cardboard sheets, to cover a 100 m<sup>2</sup> floor area. Using a paint brush or roller, the solution is applied as stripes or spots onto pieces of timber, or sheets of cardboard or fabric which are hung or attached in areas where

flies tend to aggregate. These areas may be walls, on surfaces of pillars and around windows or other structures. Treated boards should be placed in as many places as possible where flies tend to congregate (without exceeding 200g of QuickBayt per 100 m<sup>2</sup> floor/ground space).

##### Cleaning up:

DO NOT discharge any waste product or washing water from treatment equipment to drains.

Equipment used in connection with the paint-on application (e.g. brushes, roller) must not be cleaned after use. All contaminated equipment may be re-used without cleaning (if possible) and always be disposed of according to local regulations, (with no discharge into waste water).

##### Duration of effect:

QuickBayt can provide long lasting housefly control in rural hygiene and waste management facilities. However, this is likely to be best achieved if the treatment is complemented by good hygiene practice. Noticeable reduction of house fly numbers can be observed from day 1 and house fly control can be expected ≥ 3 weeks and can last up to 8 weeks under standard conditions. QuickBayt lasts for up to 8 weeks. The minimum retreatment period is 21 days. QuickBayt should not be used more than 8 times in any 12 month period.

##### SAFETY PRECAUTIONS

UK Only: The COSHH (Control of Substances Hazardous to Health) Regulation 2002 (as amended) apply to the use of this product at work. Wear protective chemical resistant gloves when handling the product. AVOID EXCESSIVE CONTAMINATION OF OVERALLS. Do not apply product to treated supports, pieces of timber, sheets of cardboard or fabric which will be placed in areas subject to washing/slaicing. DO NOT USE food, feed or water could become contaminated. Do not apply the biocidal product directly onto surfaces (e.g. walls) in the building. Do not clean the treated areas (cardboards etc...). Remove all pieces of timber, sheets of cardboard, hang-boards, fabrics and bait stations which are treated with the biocidal product before pre-cleaning and/or disinfectant events in animal housing/shelters. PREVENT ACCESS TO BAIT by children and animals. WASH HANDS AND EXPOSED SKIN before meals and after use. DO NOT touch treated surfaces until dry.

##### ENVIRONMENTAL PROTECTION:

This product contains Imidacloprid, which is toxic to bees. Upon treatment with bait, the area should be made as inaccessible to birds as possible, by closing doors/windows and/or using proofing measures to keep birds out of the area. Do not allow to get into surface water, drains and ground water. If the product is released onto soil. Methods for cleaning up: Use mechanical handling equipment. Clean contaminated floors and objects thoroughly, observing environmental regulations.

##### Resistance Management:

This product should not be used continuously against houseflies in intensive or controlled environment animal houses as this could cause failure due to insecticide resistance. Imidacloprid is the active component of QuickBayt, which belongs to the class of neonicotinoid insecticides having a different mode of action to other classes of insecticide, such as pyrethroids or organophosphates. The use pattern as paint-on bait ensures that most of the animal house is not treated thereby reducing the likelihood of contacting a sub lethal deposit. Nevertheless, to minimize the chances of resistance developing in the future, it is advisable to avoid using QuickBayt, exclusively and continuously as the sole agent for fly control. QuickBayt should be used as one component of an integrated pest management program which features products from alternative chemical classes and non-chemical means (e.g. fly screens, sanitary practices).

##### STORAGE:

Store in a dry, cool and well ventilated place. Store in original container. Keep containers tightly closed. Store in a place accessible by authorised persons only. Store away from direct sunlight. Keep away from food, drink and animal feedingstuffs. KEEP IN A SAFE PLACE. KEEP out of reach of children. This product has a shelf life of 3 years.

##### DISPOSAL OF PRODUCT AND PACKAGING:

Keep in suitable, closed containers for disposal. Remains of the applied product (e.g. empty packages, exhausted card boards, exhausted treated supports) and dead flies must be disposed of according to local waste regulations. This material and its container must be disposed of in a safe way prior to any washing/slaicing taking place.

##### FIRST AID

**General:** Move out of dangerous area. Place and transport victim in stable position (lying sideways). Remove contaminated clothing immediately and dispose of safely. **Skin Contact:** Wash off thoroughly with plenty of soap and water, if available with polyethyleneglycol 400, subsequently rinse with water.

**Eye Contact:** Rinse immediately with plenty of water, also under the eyelids, for at least 15 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Get medical attention if irritation develops and persists.

**Ingestion:** Rinse mouth. Do not give fluids or induce vomiting. Call physician or poison control centre immediately.

**Treatment:** Treat symptomatically.

**Monitor:** Respiratory and cardiac functions.

**There is no specific antidote.**

**Poison Information for Ireland: For information or to report a poisoning incident contact The National Poisons Information Centre, Beaumont Hospital, Dublin (01-8092166), retain the label for reference.**

GB05956136J - A5 - ARTICLE 05814250

Authorisation holder and marketing company:

|                             |                             |
|-----------------------------|-----------------------------|
| <b>United Kingdom</b>       | <b>Ireland</b>              |
| Bayer CropScience Ltd       | Bayer Limited               |
| 230 Cambridge Science Park, | The Atrium,                 |
| Milton Road, Cambridge,     | Blackthorn Road, Sandryford |
| CB4 0WB                     | Dublin 18                   |
| United Kingdom              | Ireland                     |

For 24 hour emergency information contact: Bayer CropScience Ltd. Telephone: 00800 1020 3333

Further information is available from: 00800 12149451

[www.es.bayer.co.uk](http://www.es.bayer.co.uk) for SDS & larger label.

**Use biocides safely and sustainably. Always read the label and product information before use.**

**It is illegal to use this product for uses or in a manner other than that prescribed on this label.**

Ⓞ QuickBayt is a Registered Trade Mark of Bayer.

Ⓞ Bitrex is a Registered Trade Mark of Macfarlan Smith, Edinburgh.

Ⓞ Copyright of Bayer

Date of expiry: see the pack. Batch number: see the pack.

**Bayer**



# QuickBayt®



# QuickBayt®

## INSECTICIDE


- ✓ For use only as an insecticide (PT18).
- ✓ For use only by professional operators.
- ✓ For indoor use only.



Paint on fly control



Safety information  
**QUICKBAYT**  
 Contains 0.5% w/w Imidacloprid & 0.1% Muscalure.



**WARNING**  
**Very toxic to aquatic life with long lasting effects.**  
 Collect spillage.  
 Avoid release to the environment.  
 Contains Cis-tricos-9-ene. May produce and allergic reaction.  
 Dispose of contents / container to a licensed hazardous-waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste.

To avoid risks to human health and the environment, comply with the instructions for use.

UK-2017-1090, IE/BPA 70269

4 x 2 kg e

## INSECTICIDE


- ✓ For use only as an insecticide (PT18).
- ✓ For use only by professional operators.
- ✓ For indoor use only.



Paint on fly control



Safety information  
**QUICKBAYT**  
 Contains 0.5% w/w Imidacloprid & 0.1% Muscalure.



**WARNING**  
**Very toxic to aquatic life with long lasting effects.**  
 Collect spillage.  
 Avoid release to the environment.  
 Contains Cis-tricos-9-ene. May produce and allergic reaction.  
 Dispose of contents / container to a licensed hazardous-waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste.

To avoid risks to human health and the environment, comply with the instructions for use.

UK-2017-1090, IE/BPA 70269

4 x 2 kg e

GB05986728J - A5

Authorisation holder and marketing company:  
**United Kingdom**  
 Bayer CropScience Ltd  
 230 Cambridge Science Park,  
 Milton Road,  
 Cambridge,  
 CB4 0WB  
 United Kingdom

**Ireland**  
 Bayer Limited  
 The Atrium,  
 Blackthorn Road,  
 Sandymount,  
 Dublin 18  
 Ireland

Bayer

For 24 hour emergency information contact: Bayer CropScience Ltd. Telephone: 00800 1020 3333  
 Further information is available from: 00800 12149451  
[www.es.bayer.co.uk](http://www.es.bayer.co.uk) for SDS & larger label.

Use biocides safely and sustainably. Always read the label and product information before use. It is illegal to use this product for uses or in a manner other than that prescribed on this label.

© QuickBayt is a Registered Trade Mark of Bayer.  
 © Bitrex is a Registered Trade Mark of Macfarlan Smith, Edinburgh.  
 © Copyright of Bayer

Date of expiry: see the pack. Batch number: see the pack.